

Razón personas/perros obtenidos a partir de estudios detallados en diversos entornos urbanos y rurales de todo el mundo, que se pueden utilizar como información preliminar sobre la cantidad de perros con dueño para la de planificación y adquisición de vacunas (por ejemplo, en el caso de campañas con que deben iniciarse con urgencia).

País	Entorno	Personas por perro	Fuente	Notas
África – todos	Urbano	21.20	Knobel et al. 2005	Relación promedio para toda la región
África – todos	Rural	7.40	Knobel et al. 2005	Relación promedio para toda la región
Tanzania	Rural: agrícola-pastoral	7.26	Kaare et al. 2009	
Tanzania	Rural: pastoral	7.55	Kaare et al. 2009	
Tanzania	Rural: interior	7.60	Knobel et al. 2008	
Tanzania	Rural: costero	10.80	Knobel et al. 2008	
Tanzania	Urbano: interior	14.40	Knobel et al. 2008	
Tanzania	Urbano: costero	27.20	Knobel et al. 2008	
Nigeria	Urbano	4.10	El-Yuguda et al. 2007	
Nigeria	Rural	3.20	El-Yuguda et al. 2007	
Kenia	Rural, periurbano	8.00	Kitala et al. 2001	
Madagascar	Urbano	4.50	Ratsitorahina et al. 2009	
Zambia	Urbano	45.00	De Balogh et al. 1993	
Zambia	Rural	6.70	De Balogh et al. 1993	
Sudáfrica	Rural	11.10	Rautenbach et al. 1991	
Zimbabue	Urbano	16.00	Brooks 1990	
Zimbabue	Rural	4.50	Brooks 1990	
Chad	Urbano	21.50	Mindekem et al. 2005	
Asia – todos	Urbano	7,50	Knobel et al. 2005	Relación promedio para toda la región
Asia – todos	Rural	14.30	Knobel et al. 2005	Relación promedio para toda la región
India		36.00	Sudarshan et al. 2006	
Tailandia		4.60	Kongkaew et al. 2004	
Filipinas		3.80	Robinson et al. 1996	
China – todos	Urbano y rural	48.30	Knobel et al. 2005	Relación promedio para toda la región
México	Urbano	3.40 – 4.30	Fishbein et al. 1992; Flores-Ibarra & Estrella-Valenzuela 2004	
Bolivia	Urbano	4.60	Suzuki et al. 2008	
Brasil	Urbano	4.00	Alves et al. 2005	
Ecuador	Urbano	7.60	Beran & Frith 1988	
Norteamérica y Europa		6.00 – 10.00	Wandeler et al. 1988	

Referencias del cuadro:

Alves MC, Matos MR, Reichmann Mde L, Dominiguez MH (2005). Estimation of the dog and cat population in the State of São Paulo. Rev Saude Publica 39: 891 – 897.

Beran GW, Frith M (1988). Domestic animal rabies control: an overview. Rev Infect Dis 10 Suppl 4:S672 – 677.

Brooks R (1990). Survey of the dog population of Zimbabwe and its level of rabies vaccination. Vet Rec 127: 592 – 596.

- De Balogh KK, Wandeler AI, Meslin FX (1993). A dog ecology study in an urban and a semi-rural area of Zambia. *Onderstepoort J Vet Res* 60: 437 – 443.
- El-Yuguda AD, Baba AA, Baba SSA (2007). Dog population structure and cases of rabies among dog bite victims in urban and rural areas of Borno State, Nigeria. *Trop Vet* 25: 34 – 40.
- Fishbein DB, Frontini MG, Dobbins JG, Flores Collins E, Quiroz Huerta G, Gamez Rodriguez JJ, Woo-Ming B, Garza Ramos J, Belotto AJ, Balderas Torres JM, et al. (1992). Prevention of canine rabies in rural Mexico: an epidemiologic study of vaccination campaigns. *Am J Trop Med Hyg* 47: 317 – 327.
- Flores-Ibarra M, Estrella-Valenzuela G (2004). Canine ecology and socioeconomic factors associated with dogs unvaccinated against rabies in a Mexican city across the US-Mexico border. *Prev Vet Med* 62: 79 – 87.
- Kaare M, Lembo T, Hampson K, Ernest E, Estes A, Mentzel C, Cleaveland S (2009). Rabies control in rural Africa: evaluating strategies for effective domestic dog vaccination. *Vaccine* 27: 152 – 160.
- Kitala P, McDermott J, Kyule M, Gathuma J, Perry B, Wandeler A (2001). Dog ecology and demography information to support the planning of rabies control in Machakos District, Kenya. *Acta Trop* 78: 217 – 230.
- Knobel DL, Cleaveland S, Coleman PG, Fèvre EM, Meltzer MI, Miranda ME, Shaw A, Zinsstag J, Meslin FX (2005). Re-evaluating the burden of rabies in Africa and Asia. *Bull World Health Organ* 83: 360 – 368.
- Knobel DL, Laurenson MK, Kazwala RR, Boden LA, Cleaveland S (2008). A cross-sectional study of factors associated with dog ownership in Tanzania. *BMC Vet Res* 4: 5.
- Kongkaew W, Coleman P, Pfeiffer DU, Antarasena C, Thiptara A. (2004). Vaccination coverage and epidemiological parameters of the owned-dog population in Thungsong District, Thailand. *Prev Vet Med* 65: 105 – 115.
- Mindekem R, Kayali U, Yemadji N, Ndoutamia AG, Zinsstag J (2005). Impact of canine demography on rabies transmission in N'djamena, Chad. *Med Trop (Mars)* 65: 53 – 58.
- Ratsitorahina M, Rasambainarivo JH, Raharimanana S, Rakotonandrasana H, Andriamiarisoa MP, Rakalomanana FA, Richard V (2009). Dog ecology and demography in Antananarivo, 2007. *BMC Vet Res* 5: 21.
- Rautenbach GH, Boomker J, de Villiers IL (1991). A descriptive study of the canine population in a rural town in southern Africa. *J S Afr Vet Assoc* 62: 158 – 162.
- Robinson LE, Miranda ME, Miranda NL, Childs JE (1996). Evaluation of a canine rabies vaccination campaign and characterization of owned-dog populations in the Philippines. *Southeast Asian J Trop Med Public Health* 27: 250 – 256.
- Sudarshan MK, Mahendra BJ, Madhusudana SN, Ashwoath Narayana DH, Rahman A, Rao NS, X-Meslin F, Lobo D, Ravikumar K, Gangaboraiah (2006). An epidemiological study of animal bites in India: results of a WHO sponsored national multi-centric rabies surveys. *J Commun Dis* 38: 32 – 39.
- Suzuki K, Pereira JA, Frías LA, López R, Mutinelli LE, Pons ER (2008). Rabies-vaccination coverage and profiles of the owned-dog population in Santa Cruz de la Sierra, Bolivia. *Zoonoses Public Health* 55: 177 – 183.
- Wandeler AI, Budde A, Capt S, Kappeler A, Matter H (1988). Dog ecology and dog rabies control. *Rev Infect Dis* 10 Suppl 4: S684 – 688.